

REMOTE SENSING APPLICATIONS IN FORESTRY

CLOUD COVER INTERFERENCE WITH REMOTE SENSING
OF FORESTED AREAS FROM EARTH-ORBITAL
AND LOWER ALTITUDES

R-09-038-002

By
Jerry D. Lent

School of Forestry
University of California

Annual Progress Report

30 September, 1966

A report of research performed under the auspices of the
FORESTRY REMOTE SENSING LABORATORY,
BERKELEY, CALIFORNIA—

A Coordination Facility Administered Jointly By

The Pacific Southwest Forest and Range Experiment Station of the
Forest Service, U.S. Department of Agriculture and by the
School of Forestry, University of California

For

NATURAL RESOURCES PROGRAM
OFFICE OF SPACE SCIENCES AND APPLICATIONS
NATIONAL AERONAUTICS AND SPACE
ADMINISTRATION

A B S T R A C T

Most remote sensing devices do not adequately image the Earth's surface when separated from it by cloud cover. Since few weather stations are located in forested areas, little has been known about the potential limitations of cloud cover for the remoted sensing of forested areas. This study analyzes the cloud cover problem for selected forested areas in California, including the area in which the NASA Forestry Test Site (the Bucks Lake Test Site) is located.

Cloud cover information from several selected meteorological stations in California was interpreted in order to determine its influence on remote sensing applications over forested areas. The question of predicting percent cloud cover at various altitudes was examined by relating recorded ground station observations with representative Earth-orbital cloud imagery. A comprehensive bibliography of recent research on cloud cover conditions, including satellite applications, is included.

AUTHOR

ACKNOWLEDGMENTS

This research was performed under the sponsorship and financial assistance of the National Aeronautics and Space Administration for the Manned Earth Orbital Experiment Program in Agriculture/Forestry (Contract number R-09-038-002 ---Natural Resources Program).

The author would like to thank Mr. Raymond Gosden for his diligent assistance in assembling the bibliographic citations included in this report. Appreciation is also directed to Miss Evelyn Hall who assisted in the data card keypunching.

TABLE OF CONTENTS

ABSTRACT	i
ACKNOWLEDGMENTS	ii
INTRODUCTION	1
JUSTIFICATION	3
METHODS AND PROCEDURES	4
DISCUSSION OF RESULTS	6
WEATHER STATION INDEX MAP	8
ILLUSTRATIONS	9
APPENDIX (MONTHLY SUMMARIES)	21
APPENDIX (BIBLIOGRAPHY)	29

CLOUD COVER INTERFERENCE WITH REMOTE SENSING OF FORESTED AREAS FROM
EARTH-ORBITAL AND LOWER ALTITUDES

by

Jerry D. Lent

INTRODUCTION

The rate at which remote sensing applications have increased in recent years has highlighted certain limitations now coming under closer examination due to the need for rapid, precise interpretation of data that can be assembled and efficiently disseminated to the scientific world. This study is concerned with one of the major sources of remote sensing limitations - the very atmosphere through which information is sought. More specifically, this study is intended to examine the need for predicting cloudiness conditions over forested areas and the accuracy with which these predictions can be made.

Cloud-free skies provide both the viewer, and most remote sensing equipment, with an unobstructed view of the terrain. Unfortunately, the Earth's surface is 50 to 55 percent obscured by clouds (on the average) at any specified time. This would present an harassing problem for scientists requiring "the best" terrain imagery were it not for the following generalities:

- a) Summer and fall seasons are less cloudy than winter and spring seasons (for the Northern Hemisphere).
- b) Arctic and equatorial regions tend to be less cloudy than temperate regions.

- c) Diurnal fluctuations in cloud amounts exist.
- d) Water bodies tend to be less cloudy than land surfaces. (See figures 1 and 9a-o.)

Planning a flight mission, whether it be low altitude or Earth-orbital, requires accurate weather information in order to predict success in obtaining certain types of terrain imagery. Cloudiness over forested areas will usually delay a flight mission designed to inventory the wildland resources. In the case of forest fire detection and disease and insect control, timing of imagery is of the utmost importance. Remote sensing systems relying on conventional imagery (i.e., panchromatic and color photography) are hindered by clouds even when cloudiness is as low as two to four-tenths skycover. When such conditions prevail, flight missions must be rescheduled for a more favorable situation (though not necessarily a more favorable time) to retrieve optimum information. (See figs. 2 and 3.)

A brief examination of the electromagnetic spectrum reveals the requirements necessary to overcome or at least reduce atmospheric interference in obtaining specified imagery. Each film and/or recorder has a specified region of sensitivity and, with it, a compromise between resolution and attenuation due to atmosphere. Orthochromatic film is sensitive at short wavelengths and consequently has poor penetrating power through atmospheric haze. Panchromatic films extend to longer wavelengths but corrective filters are required to reduce the problem of haze; conventional color films have sensitivities beyond the panchromatic range (i.e., at longer wavelengths) and thus possess greater haze penetrating

ability. As illustrated in figure 4, however, severe atmospheric haze ("smog") can seriously deteriorate the image. This emphasizes the fact that atmospheric particle size is directly related to the ease with which remote sensing can penetrate dust, smoke, and clouds. Figure 5 reveals a radar image taken with considerably longer wavelengths that essentially "see through" particles of the size that contribute to a cloud formation. Radar imagery also is capable of registering detail at night, as well as during the day, should the need arise. Since cloud particles are somewhat larger than smoke particles, this suggests that a sensing device for detecting forest fires (even under obscured skies) using the same wavelengths as are used in radar sensors, might prove superior to conventional methods of detection now in operation. However, at these radar wavelengths greater success for fire detection might be obtained through use of a "passive" microwave sensor rather than an "active" radar system.

Greater detail and image resolution is possible using panchromatic type films (see figure 6) than with radar or passive microwave sensors. For general, gross surveys and special applications, however, radar or passive microwave sensors may be more desirable whenever clouds are a problem.

JUSTIFICATION

The necessity for predicting cloudiness over either a specific region or a very large portion of the Earth has far-reaching implications. From a forestry standpoint, these implications can be expressed in terms of the ease with which land-use surveys are made, timber species and quantities delineated, management decisions formulated, forest protection practices intensified, and many other forestry applications developed.

METHODS AND PROCEDURES

In conducting this study of cloud amounts over forested areas, it was decided that California would provide suitable regions to investigate for the following reasons:

- a) the diversity of forested and agricultural conditions
- b) the well distributed weather station coverage
- c) the complexity of meteorological conditions, and
- d) the presence of NASA test sites for which cloudiness is an important concern in flight mission planning.

Cloudiness information for eight stations was obtained from the Weather Records Center, Asheville, North Carolina. This information consisted of magnetic tapes and punched cards. In addition, representative TIROS imagery was obtained as an aid in interpreting data on cloud amounts. (Refer to figs. 7-9.) The Reno weather station was included to indicate how cloudiness changes from the crest of the California Sierra Nevada Range to the Great Basin Plateau. This information is included in the "Results" section where Blue Canyon cloudiness is compared with that of the Reno station.

Where possible, each station was analyzed for a fifteen year period. Monthly summaries were made first and variations of cloudiness tested so that 95 and 90 percent confidence bands, respectively, were computed. Blue Canyon and Reno data are attached herein. Simply, the method of computing these confidence bands is:

<u>Confidence level</u>	<u>Confidence band</u>
95%	monthly average \pm 1.96 (standard deviation of data in population)
90%	monthly average \pm 1.64 (standard deviation of data in population)

The ninety-five percent (upper) confidence level was chosen because it was felt that, owing to the nature of weather data, a greater confidence requirement would result in too broad an interval for realistic cloudiness conditions in any given area. The ninety percent (lower) confidence level was chosen because it was felt that little significance could be ascribed to cloud-cover predictions based on a degree of confidence lower than this.

By applying the foregoing simple analysis to the weather stations incorporated in this study (see map, page 8), it was then possible to summarize in tabular form the levels of confidence associated with the specified time interval of cloudiness data. Cloudiness data for specified altitudes and times of day were available, but cursory examination of these suggested that less than meaningful information would result from the method selected for analysis. Consequently, monthly summaries are presented herein without regard to differences in cloud altitudes. It should be emphasized that long term records of cloudiness data are currently undergoing adaption to EDP files so that records longer than those employed in this study can be analyzed quickly in the future.

The generalization regarding cloudiness conditions over a forested region--which might result from a few, widely scattered weather station skycover reports--may prove entirely adequate for certain objectives; but, for the most part, these generalizations must be loosely interpreted. Meteorological variables have been shown to be complex in nature, and considerable "homogeneity" of terrain is required before wide-ranging extrapolations between observing stations are made; in mountainous, forested regions these extrapolations are confounded by a multitude of variables which make predictions of clouds a generalization at best.

DISCUSSION OF RESULTS

Two viewpoints of analysis commonly held by meteorologists--numerical analysis and synoptic analysis--have been incorporated in this study as an aid to deriving meaningful results.


The sparsity of weather stations so situated as to record cloud cover in remote forested areas presented problems, in this study, of extrapolating data between stations. A similar difficulty would likely exist with respect to most other forested areas of the world. In the absence of an adequate network of weather stations, the problems associated with predicting cloud amounts of a specified forested region are very complex. It is readily seen by noting figure 1 that present weather stations recording cloud cover are located in well established sites. The forested regions of the southeast United States, in fact, are sufficiently homogeneous in land features that a closer interpretation of data (vis., cloudiness at various altitudes and times of day) is possible with reliability for success greatly enhanced. Along the Pacific Coast Range and Rocky Mountain Range, however, this homogeneity deteriorates into a myriad of terrain characteristics such that forested regions remain relatively remote even by present standards. Many meteorological variables can contribute to cloudiness and the interplay of these variables is not completely understood at the present time. Promising factors for additional study include upper air wind patterns that contribute to changing weather conditions, precipitation records, and radiation measurements.

The use of weather satellite imagery is a proven application for instantaneous weather surveys throughout the world. The latest Nimbus series of weather satellites provides the Weather Bureau with the best available source of rapid world-wide weather data now known. Nephographs showing cloud

type and distribution are drawn up, and when compared with long-term ground observations, show broad agreement. It should be mentioned that the resolution of such imagery is such that little extrapolation beyond meteorological aspects is commonly made. To the meteorologist, clouds become a "target" rather than an obstruction.

The following tables list summaries of cloudiness over a period of years for the stations included in this study: Blue Canyon, located at the crest of the central Sierra Nevada, and Reno, Nevada, some thirty miles away are especially interesting to compare. The Reno station is broken down into nighttime hours and daytime hours and it is seen that clouds are less frequent during the nighttime hours on the average. In comparing the two tables the difference between nighttime and daytime cloudiness for the Reno station must be considered. The most cloud-free periods for these two stations are July and August for Blue Canyon (averaging close to twenty cloud-free days a month) and August and September for Reno (averaging fifteen cloud-free days a month). Timing of ground observations is at six-hour intervals. Afternoon build-up in the Sierra Nevada during these summer months somewhat confuses the problem of recording cloud cover.

In conclusion, this author feels that useful generalizations presently can be made regarding the weather situation likely to exist at given times over a given forested area. When it appears probable from these generalizations that cloud cover will obscure forested areas of interest from aerial view, radar or passive microwave imagery may still provide very useful information despite its present limited resolution characteristics. Continuation of basic research of the type suggested and reported herein, including extensive use of data available from weather reconnaissance satellites, should steadily improve the accuracy with which these vital weather predictions can be made.



The above map of California depicts the location of weather stations used in assembling data on cloudiness for this study; the Bucks Lake Test Site is some 50 miles north of Blue Canyon (1); Pisgah Crater is only 11 miles from the Daggett station (5); the Davis agricultural Test Site is 20 miles west of the Sacramento station (7); the San Pablo Reservoir Test Area is only a few miles from the Oakland station.




Figure 1: (Ref. No. 22, Bibliography Section) Long term cloudiness charts for the United States. These diagrams are constructed from weather station data listing "skycover" as a percent of total cloudiness. Weather stations denoting cloudiness are indicated by the "dots" shown above. The generalities regarding cloudiness as depicted by charts like those shown here are sufficiently detailed for most purposes.



a) Ekta Aero Infrared (oblique-30000ft. above terrain)



b) Ekta Aero Infrared (vertical-scale: 1/20,000)


Figure 2: Proper selection of film type to meet a mission's objective can often be disrupted by atmospheric interference. In the above example illustrations, it is seen that clouds have obscured terrain features sufficiently to deplete information which might be obtained. The "bluish" cast in (a) above is due to the great depth of atmospheric penetration required for this photographic angle. In (b) the San Pablo Reservoir Test Site is shown which is subjected to dense fog banks at certain times of the year.



a) Kodachrome



b) Ektachrome



c) Ekta Aero Infrared

Figure 3: Many types of atmospheric interference are known. The above example illustrations are pertinent to several California NASA Test Sites. (a) illustrates a phenomenon especially peculiar to the Pisgah Crater Site -- high speed aircraft contrails from nearby airbases. (b) illustrates a fog bank near the San Pablo Reservoir Test Site filmed through emulsions sensitive to different parts of the Electromagnetic spectrum.



Figure 4: Severe atmospheric interference is becoming more common in low-land urban and suburban regions of the United States. The above example ground illustration shows a Southern California vineyard subjected to moderate "smog" conditions exposed through black and white infrared film. While this condition is relatively rare in regional forested areas, it is noted that local sawmill activity can generate conditions similar to those above under inversion atmospheres.




Figure 5: The application of remote sensing employing longer wavelengths than conventional imagery (i.e., panchromatic and near-infrared) is gaining support from many scientific disciplines. The example illustration shown here depicts a radar exposure of the Bucks Lake Test Site (California). Proper selection of radar or passive microwave sensors operating in the longer wavelengths prove invaluable aids to forestry applications when clouds obscure the terrain or when detection of forest fires is the objective. The reduced resolution inherent in present radar systems must be weighed against the ability to obtain imagery at a time when it is needed, irrespective of atmospheric interference. Texture and tone differences aid the interpreter in distinguishing brush from timber; large water bodies are readily distinguished by tone; even burned brushfields can be seen as darkened areas in the circled area above. (The square outline shows approximate area seen on Figure 6).




Figure 6: A portion of the area seen in Figure 5 is shown here in conventional panchromatic film. Greater detail is possible, but even here there is sufficient cloud obstruction to hinder accurate delineation of the waterline. Cloud shadows also reduce interpretability (see Figure 11).




Figure 7: Example of TIROS imagery and accompanying reference data. Taped and Direct TV transmission are photographed on 35 mm black-and-white film.







Figure 8: Earth-orbital image of Grand Bahama Reef obtained from an early Gemini mission showing resolution and color advantages.




9a)


9b)




9c)




9d)




9e)




9f)




9i)



9l)




9h)




9k)




9g)




9j)



9m)



9n)



9o)





Figure 10: High altitude oblique and vertical images. Left illustration shows attenuating atmosphere on horizon and cloud development near mountain crests. Right illustration is developed in such a way to bring out cloud features, not terrain features, and reveals suitability of accurate height measurements using photogrammetric principles.





Figure 11: The above two illustrations point out the "double interference" usually associated with clouds during conventional photographic missions. Not only are the clouds obscuring important terrain detail, but their shadows are similarly obscuring the terrain. Exposure to compensate for cloud normally results in poorly degraded imagery where no clouds exist (see Figure 10).

Table 1. Blue Canyon, California
(length of record: 5 years)

Cloudiness data computed from 0 to 10 (clear to 100% sky cover)

Confidence Bands for Cloudiness Data*

<u>Month</u>	<u>Mean</u>	<u>90%</u>	<u>95%</u>
January	5.4	9.0 - 1.8	9.7 - 1.1
February	6.4	9.2 - 3.6	9.7 - 3.1
March	6.7	8.0 - 5.4	8.3 - 5.1
April	6.6	9.8 - 3.4	10.0 - 2.8
May	6.0	8.0 - 4.0	8.4 - 3.6
June	3.6	5.7 - 1.5	6.1 - 1.1
July	0.8	1.7 - 0.0	1.8 - 0.0
August	2.4	4.4 - 0.4	4.8 - 0.0
September	2.4	4.0 - 0.8	4.4 - 0.4
October	3.9	6.1 - 1.7	6.5 - 1.3
November	6.3	7.5 - 5.1	7.7 - 4.9
December	6.3	8.4 - 4.1	8.8 - 3.8

*Interpreted as the percent that the true cloudiness condition will lie within the band specified.

Table 2 - Reno, Nevada

(length of record: 8 years)

Cloudiness data computed from 0 to 10 (clear to 100% skycover)

Confidence Bands for Cloudiness Dates*

Month	Mean	<u>Daylight hours</u>		Mean	<u>Nighttime hours</u>	
		<u>90%</u>	<u>95%</u>		<u>90%</u>	<u>95%</u>
J	6.1	9.1-3.1	9.6-2.6	5.4	7.7-3.1	8.6-2.2
F	6.2	9.5-2.9	10.0-2.3	5.6	8.7-2.5	9.3-1.9
M	6.0	7.4-4.6	7.7-4.3	5.4	7.1-3.7	7.4-3.4
A	5.6	7.1-4.1	7.3-3.9	4.9	6.5-3.3	6.8-3.0
M	5.2	7.0-3.4	7.4-3.0	4.8	6.6-3.0	6.9-2.7
J	3.6	5.4-1.8	5.8-1.4	3.4	5.1-1.7	5.5-1.3
J	2.3	4.0-0.6	4.3-0.3	2.1	3.7-0.5	4.0-0.2
A	2.5	4.1-0.9	4.2-0.8	2.3	4.1-0.5	4.4-0.2
S	2.5	4.2-0.8	4.6-0.4	2.1	3.6-0.6	3.9-0.3
O	3.8	5.5-2.1	5.8-1.8	3.0	4.5-1.5	4.8-1.2
N	5.5	8.2-2.8	8.7-2.3	4.7	7.1-2.3	7.6-1.8
D	5.8	10.0-1.4	10.0-0.5	4.9	6.6-3.2	6.9-2.9

*Interpreted as the percent that the true cloudiness condition will be within the band specified.

Table 3. Arcata NAAS, California
(length of record: 15 years)

Cloudiness data computed from 0 to 10 (clear to 100% sky cover)

Confidence Bands for Cloudiness Data*

<u>Month</u>	<u>Mean</u>	<u>90%</u>	<u>95%</u>
January	6.5	8.0 - 5.0	8.3 - 4.7
February	6.2	8.1 - 4.3	8.9 - 3.5
March	6.1	8.1 - 4.1	9.0 - 3.2
April	5.9	8.2 - 3.6	8.5 - 3.3
May	5.6	7.6 - 3.6	7.9 - 3.3
June	5.4	7.3 - 3.5	7.7 - 3.1
July	5.0	7.0 - 3.0	7.2 - 2.8
August	4.7	6.9 - 2.5	7.2 - 2.2
September	4.9	7.4 - 2.4	7.9 - 1.9
October	5.1	8.4 - 1.8	8.7 - 1.5
November	5.8	9.2 - 2.4	9.4 - 2.2
December	6.6	8.5 - 4.7	8.7 - 4.5

*Interpreted as the percent that the true cloudiness condition will lie within the band specified.

Table 4 - Bishop, California
(length of record: 15 yrs)

Cloudiness data computed from 0 to 10 (clear to 100% skycover)

Confidence Bands for Cloudiness Data*

<u>Month</u>	<u>Mean</u>	<u>90%</u>	<u>95%</u>
January	4.3	7.5 - 1.1	7.8 - 0.8
February	4.2	7.0 - 1.4	7.2 - 1.2
March	4.0	5.5 - 2.5	5.7 - 2.3
April	3.4	4.8 - 2.0	4.9 - 1.9
May	2.3	4.3 - 0.3	4.4 - 0.2
June	2.0	3.7 - 0.3	4.1 - 0.0
July	1.7	2.5 - 0.8	2.7 - 0.6
August	2.0	3.2 - 0.8	3.3 - 0.7
September	2.1	4.2 - 0.0	4.4 - 0.0
October	2.7	5.3 - 0.1	5.5 - 0.0
November	3.4	5.9 - 0.9	6.2 - 0.6
December	4.5	7.5 - 1.5	7.8 - 1.2

*Interpreted as the percent that the true cloudiness condition will lie within the band specified.

Table 5 - Daggett, California
(length of record: 15 yrs)

Cloudiness data computed from 0 to 10 (clear to 100% skycover)

Confidence Bands for Cloudiness Data*

<u>Month</u>	<u>Mean</u>	<u>90%</u>	<u>95%</u>
January	3.0	3.3 - 2.7	3.4 - 2.6
February	3.1	3.6 - 2.6	3.8 - 2.4
March	3.3	3.9 - 2.7	4.1 - 2.5
April	2.5	3.1 - 1.9	3.2 - 1.8
May	2.0	2.4 - 1.6	2.5 - 1.5
June	1.6	1.9 - 1.3	2.0 - 1.4
July	1.5	2.0 - 1.0	2.2 - 0.8
August	1.8	2.1 - 1.5	2.3 - 1.3
September	1.9	2.2 - 1.6	2.5 - 1.3
October	2.2	2.6 - 1.8	2.9 - 1.5
November	2.3	3.0 - 1.6	3.4 - 1.2
December	2.8	3.6 - 2.0	4.0 - 1.6

*Interpreted as the percent that the true cloudiness condition will lie within the band specified.

Table 6 - Oakland, California
(length of record: 15 yrs)

Cloudiness data computed from 0 to 10 (clear to 100% skycover)

Confidence Bands for Cloudiness Data*

<u>Month</u>	<u>Mean</u>	<u>90%</u>	<u>95%</u>
January	5.2	8.0 - 2.4	8.2 - 2.2
February	5.0	8.3 - 1.7	8.5 - 1.5
March	4.9	7.4 - 2.4	7.7 - 2.1
April	4.3	6.6 - 2.0	6.8 - 1.8
May	3.3	6.4 - 0.2	6.5 - 0.1
June	3.0	5.8 - 0.2	6.0 - 0.0
July	2.8	5.5 - 0.1	5.7 - 0.0
August	3.0	5.6 - 0.4	5.8 - 0.2
September	3.4	6.2 - 0.6	6.4 - 0.4
October	3.6	6.5 - 0.7	6.7 - 0.5
November	3.7	7.2 - 0.2	7.5 - 0.0
December	5.0	7.7 - 2.3	7.9 - 2.1

*Interpreted as the percent that the true cloudiness condition will lie within the band specifies.

Table 7 - Sacramento, California
(length of record: 15 yrs)

Cloudiness data computed from 0 to 10 (clear to 100% skycover)

Confidence Bands for Cloudiness Data*

<u>Month</u>	<u>Mean</u>	<u>90%</u>	<u>95%</u>
January	4.8	7.9 - 1.7	8.0 - 1.6
February	4.0	6.6 - 1.4	6.8 - 1.2
March	3.8	6.1 - 1.5	6.2 - 1.4
April	3.7	5.6 - 1.8	5.8 - 1.6
May	3.1	4.4 - 1.9	4.5 - 1.8
June	2.6	3.6 - 1.6	3.8 - 1.4
July	1.7	2.2 - 1.2	2.2 - 1.1
August	0.8	1.2 - 0.4	1.2 - 0.4
September	1.5	2.0 - 1.0	2.1 - 0.8
October	1.8	2.8 - 0.9	2.9 - 0.7
November	2.9	4.2 - 1.6	4.4 - 1.4
December	4.4	7.2 - 1.6	7.4 - 1.4

*Interpreted as the percent that the true cloudiness condition will lie within the band specified.

Table 8 - Fresno, California
(length of record: 15 yrs)

Cloudiness data computed from 0 to 10 (clear to 100% skycover)

Confidence Bands for Cloudiness Data*

<u>Month</u>	<u>Mean</u>	<u>90%</u>	<u>95%</u>
January	3.7	6.4 - 1.0	6.6 - 0.8
February	3.9	6.4 - 1.4	6.7 - 1.1
March	3.8	5.6 - 2.0	5.8 - 1.8
April	3.1	4.4 - 1.8	4.7 - 1.5
May	2.4	3.5 - 1.4	3.5 - 1.5
June	1.8	2.6 - 1.0	2.8 - 0.8
July	1.4	1.8 - 1.0	1.8 - 0.9
August	0.8	1.1 - 0.5	1.3 - 0.4
September	1.5	2.1 - 0.9	2.4 - 0.6
October	2.3	3.4 - 1.1	3.6 - 0.9
November	3.2	5.4 - 1.0	5.8 - 0.6
December	6.0	10.0 - 1.2	10.0 - 0.4

*Interpeted as the percent that the true cloudiness condition will lie within the band specified.

BIBLIOGRAPHY

THE FOLLOWING SELECTED BIBLIOGRAPHIC CITATIONS ARE INCLUDED TO INDICATE THE LEVEL OF CONTEMPORARY INVESTIGATIONS OF CLOUDINESS AND ITS INFLUENCE ON REMOTE SENSING APPLICATIONS. THEY REVEAL MAJOR CONTRIBUTIONS, PERTINENT TO THIS PRESENT INVESTIGATION, FROM SEVERAL DISCIPLINES INCLUDING METEOROLOGICAL ASPECTS, STATISTICAL APPLICATIONS, EARTH-ORBITAL AND SUB-ORBITAL SENSING TECHNOLOGY, AND OTHERS. EACH HAS A BEARING ON THE QUESTION OF CLOUDINESS SUCH AS--PREDICTING CLOUD COVER, ANALYSIS OF DATA TRANSMITTED FROM WEATHER SATELLITES, RELATED DATA RECORDS (E.G., PRECIPITATION DATA), SHORT-TERM AND LONG-RANGE WEATHER FORECASTING, CLOUD PHOTOGRAPHY, AND METEOROLOGICAL VARIABLES AFFECTING CLOUDINESS CONDITIONS. CITATIONS DATED PRIOR TO 1958 ARE INCLUDED BECAUSE OF THEIR SPECIAL SIGNIFICANCE - THE LIST IS FAIRLY COMPLETE FROM 1958 TO THE PRESENT.

1. ABRAMOVICH, K.G.
OB USLOVIIAKH OBRAZOVANIIA OBLACHNOSTI NIZHNEGO IARUSA.
(CONDITIONS FOR THE FORMATION OF LOW LEVEL CLOUDINESS).
MOSCOW. TSENTRAL'NYI INSTITUT PROGNOZOV, TRUDY,
NO. 80, 10-41. 1959.
2. ANONYMOUS
SATELLITE AND SPACE PROBE SUMMARY--TIROS VIII.
AMERICAN METEOROLOGICAL SOCIETY, BULLETIN 45(4)
P249. 1964.
3. ANONYMOUS
TIROS I COMPLETES MISSION, TIROS II READIED.
WEATHERWISE, BOSTON. 13(4), 159-161. AUG. 1960.
4. APPLEMAN, HERBERT S. (HDQ. AIR WEATHER SERVICE SCOTT AFB)
COMPARISON OF SIMULTANEOUS AIRCRAFT AND SURFACE CLOUD
OBSERVATIONS.
JOURNAL OF APPLIED METEOROLOGY, BOSTON. 1(4),
548-551. DECEMBER 1962.
5. APPLEMAN, HERBERT S.
OCCURRENCE AND FORECASTING OF CIRROSTRATUS CLOUDS.
WORLD METEOROLOGICAL ORGANIZATION, WMO NO.
109TP47. 1961.
6. ARKING, A. (GODDARD INST FOR SPACE STUDIES, NASA N.Y.).
PERCENTAGE CLOUD COVER FROM TIROS PHOTOGRAPHY.
INT'L SPACE SCIENCE SYMPOSIUM, 4TH, WARSAW. 1963
SPACE RESEARCH, VOL. 4, PUBLISHED 1964.
7. AUBERT, EUGENE J., LUND, IVER A., THOMASELL, ALBERT JR.
SOME OBJECTIVE SIX-HOUR PREDICTIONS PREPARED BY
STATISTICAL METHODS.
8. AVERBUKH, S.K. AND POZDNIKOVA, V.A.
REZUL'TATY ISPOL'ZOVANIIA VYVODOV NEKOTORYKH ISSLE-
DOVANIIA PRI SOSTOVLENII DOLOGOSROCHNYKH PROGNOZOV
POGODY MALOI ZABLOGOVREMENNOSTI.
(THE RESULTS OF THE USE OF THE CONCLUSIONS OF CERTAIN
INVESTIGATIONS IN THE PREPARATION OF LONG-RANGE WEATHER
FORECASTS PREPARED A SHORT TIME IN ADVANCE.)
MOSCOW. TSENTRAL'NYI INSTITUT PROGNOZOV, TRUDY,
73, 100-105. 1958.
9. BALL, JOHN T.
USE OF RADIOSONDE OBSERVATIONS TO DIAGNOSE CLOUD AMOUNT
IN LAYERS.
TRAVELERS RESEARCH CENTER, INC. HARTFORD, CONN.
CONTRACT FAA/BRD-363, TECH MEMO (UNNUMBERED),
TECHNICAL PUBL. NO. 24, DECEMBER 1962.

10. BANDEEN, W.R. AND NORDBERG, W.
SUMMARY OF REFLECTED AND EMITTED RADIATION DATA
MEASURED FROM TIROSII, III, IV.
PRESENTED AT 2ND WESTERN NAT'L MEETING OF THE
AMER. GEOPHYSICAL UNION. DEC. 29, 1962.
11. BARGER, GERALD L., SHAW, ROBERT H., DALE, ROBERT F.
CHANCES OF RECEIVING SELECTED AMOUNTS OF PRECIPITATION
IN THE NORTH CENTRAL REGION OF THE UNITED STATES.
1ST RPT., AMES, IOWA. AGRIC. AND HOME ECONOMIC
EXPT. STA. IOAW ST. UNIV. JULY 1959. 277P
12. BARONI, ANDREA
LE NUBI ALTI E LA LORA PREVISIONE (HIGH CLOUDS AND
THEIR FORECASTING).
RIVISTI DI METEOROLOGIA AERONAUTICA, ROME. 19(1)
29-34. 1959.
13. BATTAU, LOUIS J.
RADAR METEOROLOGY.
CHICAGO UNIV. PRESS. 1959. 161P
14. BEST, WILLIAM H., AND PANOLSKY, H.A.
SOME APPLICATIONS OF STATISTICS TO METEOROLOGY.
COLLEGE OF MINERAL INDUSTRIES, EXTENSION SERVICES
PENN ST. COLLEGE. 1954.
15. BILANCINI, R.
LA FOTOGRAFIA DELLE NUBI (CLOUD PHOTOGRAPHY).
RIVISTI DI METEOROLOGICA AERONAUTICA, ROME. 19(2)
49-50. APRIL/JUNE 1959.
16. BOYDEN, C.J. (FORECASTING DIV., MET. OFFICE)
THE PROCESS OF FORECASTING.
MARINE OBSERVER, LONDON. 28(182), 204-206. 1958.
17. BRISTOR, C.L.
PROCESSING SATELLITE WEATHER DATA, A STATUS REPORT PT 1
1962 FALL JOINT COMPUTER CONFERENCE, AFIPS, CONF
PROCEEDINGS. VOL. 22, 11-18 (SPARTAN BOOKS,
WASHINGTON D.C.)
18. BROOKS, CHARLES F.
CLIMATIC MAPS OF NORTH AMERICA.
BLUE HILL METEOROLOGICAL OBSERVATORY, HARVARD UNIV
1936.
19. BUGAEV, V.A. AND MININE, L.S. (TSENTRAL'NYI INST. PROG.)
O GRIADOVOI STRUKTURE OBLOCHNOSTI (STRATIFICATION OF
CLOUD COVER).

METEOROLOGIJA I GIDROLOGIJA, MOSCOW. NO 5, 29-36.
MAY 1965.

20. BUNDGAARD, ROBERT C., FLETCHER, ROBERT D., SMITH, J.R.
TYPHOON-EYE CLOUD PATTERNS AS VIEWED FROM ABOVE.
WEATHERWISE, BOSTON. 12(2), 64-66. APRIL 1959.
21. BURDECKI, F.
PERSPECTIVE OF CLOUDS AS SEEN FROM SPACE VEHICLES.
SOUTH AFRICA WEATHER BUREAU, NOTOS 11(1/4), 31-45
1962.
22. CLARK, KENNETH MC R.
A NEW SET OF CLOUDINESS CHARTS FOR THE UNITED STATES.
QUARTERLY JOUR. OF THE ROYAL METEOROLOGICAL SOC.,
37(158) APRIL 1911.
23. CLAUSS, J.R.
RELATIONSHIP BETWEEN TOTAL CLOUD AMOUNT AND OBJECTIVELY
DEFINED SYNOPTIC PARAMETERS.
BERLIN. FREIE UNIVERSITAT. INST FUR METEOROLOGIE
GEOPHYSIK, METEOROLOGISCHE ABHANDLUNGEN. 32(3),
1963.
24. COLSON, DE VER
THUNDERSTORM ANALYSIS IN THE NORTHERN ROCKY MOUNTAINS.
U.S. FOREST SERVICE. INT. MTN. FOR. AND RES,
OGDEN. RESEARCH PAPER NO 49. 1957.
25. CONOVER, JOHN H.
CLOUD INTERPRETATION FROM SATELLITE ALTITUDES.
USAF. CAMBRIDGE RESEARCH LABS. RESEARCH NOTE 81,
JULY 1962.
26. CONOVER, JOHN H. AND SADDLER, JAMES C.
CLOUD PATTERNS AS SEEN FROM ALTITUDES OF 250 TO 850
MILES - PRELIMINARY RESULTS.
AMERICAN METEOROLOGICAL SOCIETY, BULLETIN 41(6),
291-297. JUNE 1960.
27. CONOVER, JOHN H.
THE IDENTIFICATION AND SIGNIFICANCE OF OROGRAPHICALLY
INDUCED CLOUDS OBSERVED BY TIROS SATELLITES.
JOURNAL OF APPLIED METEOROLOGY, BOSTON. 3(3),
226-234. JUNE 1964.
28. COUFAL, L. AND STUCHLIK, F.
OBLOCNOST V PRAZE ZA STO LET (CLOUDINESS IN PRAGUE
DURING 100 YEARS).
29. COURT, ARNOLD

THUNDERSTORM FREQUENCY IN NORTHERN CALIFORNIA.
BULLETIN, AMERICAN METEOROLOGICAL SOCIETY. 1959.

30. DADY, GUY
LA THEORIE DE L'INFORMATION DANS LES PROBLEMES
METEOROLOGIQUES.
(THE INFORMATION THEORY IN METEOROLOGICAL PROBLEMS.)
JOURNAL SCIENTIFIQUE DE LA METEOROLOGIE, PARIS.
10(39), 121-131. 1958.
31. DOOLITTLE, R.C., L. MILLER, AND I. RUFF
GEOGRAPHIC LOCATIONS OF CLOUD FEATURES.
APPENDIX I OF SOME METEOROLOGICAL RESULTS FROM
TIROS I BY STAFF. MSL, U.S. WEATHER BUREAU, IN
A REPORT BY NASA (IN PRESS, 1961.)
32. EAST AFRICAN HIGH COMMISSION, METEOROLOGICAL DEPT.
THE FREQUENCY OF TOTAL CLOUD AMOUNTS AT STATIONS IN E.
AFRICA.
(NAIROBI *) 1959.
33. ELLIOTT, F.E.
MEAN MONTHLY CLOUD COVER IN THE U.S.S.R.
G.E. ADV ELECT CENTER, CORNELL UNIVERSITY, ITHICA
TECH INFO SERIES NO. R60ELC 31 MAR 1960
34. ERICKSON, C.O. AND HUBERT, L.F.
IDENTIFICATION OF CLOUD FORMS FROM TIROS I PICTURES.
MSL REPORT NO. 7, US WEATHER BUREAU. JUNE 1961.
35. FARNHAM, D.L.
OBJECTIVE FORECASTING OF CIRRUS CLOUDINESS AT THE NAVAL
ORDNANCE TEST STATION.
U.S. NAVAL ORDNANCE TEST STATION, INYOKERN, CAL.
NAVORD REPORT NO. 6550 NOTS-TP-2249. MAY 1959.
36. FEIGEL'SON, E.M.
THE INFLUENCE OF CLOUDS ON THE ATMOSPHERIC HEAT BALANCE
IN A COLLECTION OF ARTICLES ON DYNAMIC METEOROLOG
EDITED BY I.A. KIBEL', SOVIET RESEARCH ON
GEOPHYSICS (TRANSLATED FROM RUSSION FOR AMER.
GEOPHYSICAL UNION. 1956).
37. FOX, ROY L. (USWB REG ADMIN OFF KANSAS CITY, MO.)
SUNSHINE-CLOUDINESS RELATIONSHIPS IN THE UNITED STATES.
MONTHLY WEATHER REVIEW, WASH. D.C. 89(12),
5643-5648. DECEMBER 1961.
38. FRITZ, SIGMUND
SNOW SURVEYS FROM SATELLITE PICTURES.
INT. SYMP. ON ROCKET AND SAT. MET., 1ST, WASH.,

1962 PROCEEDINGS ED. BY H. WEXLER AND T.E. CASKEY
AMSTERDAM, NORTH-HOLLAND PUB. CO. P419-421. 1963.

39. FRITZ, SIGMUND, WINSTON, J.S.
SYNOPTIC USE OF RADIATION MEASUREMENTS FROM SATELLITE
TIROS II.
REPRINTED FROM MONTHLY WEATHER REVIEW VOL. 90
P1-9. JANUARY 1962.
40. FRITZ, SIGMUND (US WEATHER BUREAU)
ON OBSERVING THE ATMOSPHERE FROM SATELLITES PT. I,
CLOUD OBSERVATION.
WEATHERWISE, BOSTON. 12(4), 139-144. AUG. 1959.
41. FUJITA, TETSUYA
TECHNIQUE FOR PRECISE ANALYSIS OF SATELLITE PHOTOGRAPHS
CHICAGO UNIV. DEPT. OF THE GEOPHYS. SCIENCES,
GRANT CWB-WBG 6, RES. PAPER NO 17, APR 1963. 53P
42. FUNKE, P.W.
PRE TROUGH WINTER PRECIPITATION FORECASTING.
USAF CAMBRIDGE RESEARCH CENTER, AIR FORCE SURVEY
IN GEOPHYSICS NO 84. 1957. 62P
43. GAL'PERIN, B.M. AND KUCHUMOVA, L.S.
O VLIIANII OBLACHNOSTI NA IZLUCHEMIE ATMOSFERY. (EFFECT
OF CLOUDINESS ON ATMOSPHERIC RADIATION.)
METEOROLOGIJA I GIDROLOGIJA, LENINGRAD. NO. 8,
19-24. AUGUST 1959.
44. GAL'PERIN, B.M.
POVTORIAEMOST' RAZLICHNOGO KOLICHESTVA OBLAKOV RAZNYKH
(FREQUENCY OF DIFFERENT AMOUNTS OF CLOUDS OF DIFFERENT
FORMS.)
LENINGRAD. GLAVNAIA GEOFIZICHESKAIA OBSERVATORIIA
161, 42-45. 1964.
45. GEORGE, J.J.
FURTHER STUDIES IN THE DEVELOPMENT OF SHORTRANGE
WEATHER PREDICTION TECHNIQUES.
EASTERN AIR LINES, INC. METEOROLOGY DEPT.
CONTR. AF19(604) 2073 FINAL REPORT JUNE 1960
46. GEORGE, J.J.
THE PREDICTION OF VERY LOW CEILINGS AND FOGS.
EASTERN AIR LINES, INC. MET. DEPT. CONTR.
AF19(604)-2073. SCIENTIFIC REPT NO 1, APR 1958.
P44-141.
47. GERGEN, JOHN L. AND HUCH, WILLIAM F.
A SYNOPTIC STUDY OF INFRARED RADIATION OVER THE U.S.

- NATURE, LONDON. 186(4723), 462-479. MAY 7, 1960
48. GIBBONS, MATHEW G. ET AL (NAVAL RADIOLOGICAL LAB S.F., CAL)
TRANSMISSION AND SCATTERING PROPERTIES OF A NEVADA
DESERT ATMOSPHERE UNDER CLOUDY CONDITIONS.
OPTICAL SOCIETY OF AMERICA, JOURNAL. 52(1),
38-43. JANUARY 1962.
49. GILLIAM, HAROLD
WEATHER OF THE SAN FRANCISCO BAY REGION.
BERKELEY U.C. PRESS 1962.
50. GOLDBERG, I.L.
NIMBUS HIGH RESOLUTION IR MEASUREMENTS.
SYMP. ON REMOTE SENSING OF THE ENVIRONMENT, 3RD.
UNIV. OF MICHIGAN, ANN ARBOR. FEB. 1965.
51. GRANT, HUGH DUNCAN
CLOUD AND WEATHER ATLAS.
NEW YORK. COWARD-MC CANN INC. 1944. 294P
52. GREAT BRITAIN, METEOROLOGICAL OFFICE
CLOUD FORMS ACCORDING TO THE MET. CLASS. DEFINITIONS
AND DESCRIPTIONS APPROVED BY THE UNT. MET. ORGAN.
WITH PHOTOGRAPHS OF CLOUDS.
LONDON H.M. STATIONERY OFFICE 1949
53. GREENFIELD, STANLEY M. AND KELLOGG, WILLIAM W.
WEATHER RECONNAISSANCE BY SATELLITES.
ASTRONAUTICS, NEW YORK. 4(1), 32-33. JAN. 1959.
54. GRAY, THOMAS I. AND SINGER, FRED S.
AN INVESTIGATION OF MET. SAT. CHARACTERISTICS FOR
CONTINUOUS EARTH COVERAGE.
USWB. MSL. LAB REPORT NO. 19 JULY '63 39 P.
55. HANEL, R.A.
DETERMINATION OF CLOUD ALTITUDE FROM A SATELLITE.
JOURNAL OF GEOPHYSICAL RESEARCH, VOL. 66, P1300.
1961.
56. HANEL, R.A. AND WARK, D.Q.
PHYSICAL MEASUREMENTS FROM METEOROLOGICAL SATELLITES.
ASTRONAUTICS AND AEROSPACE ENGIN., N.Y. 1(3),
85-88. APRIL 1963.
57. HANEL, R., J. LICHT, W. NORDBERG, R. STAMPFL, W.G. STROUD
THE SATELLITE VANGUARD II--CLOUD COVER EXPERIMENT.
IRE TRANS. ON MILITARY ELECTRONICS, MIL-4 (2 AND
3), 245-247. APRIL/JULY 1960.

58. HANEL, R.A. AND WARK, D.Q.
TIROS II RADIATION EXPT. AND ITS PHYSICAL SIGNIFICANCE.
JOURNAL, OPTICAL SOCIETY OF AMERICA. VOL. 51(12)
1394-1399. DECEMBER 1961.
59. HARRISON, HENRY T. (DIV. OF METEOROLOGY-UNITED AIRLINES)
FORECASTING SUB-MINIMUM FOG AT BALTIMORE FRIENDSHIP
AIRPORT.
UNITED AIRLINES INC. METEOROLOGY CIRCULAR NO. 45
1949.
60. HAWKINS, R.S.
ANALYSIS AND INTERPRETATION OF TIROS II INFRARED
RADIATION MEASUREMENTS.
JOUR. OF APPLIED METEOROLOGY, 3(5), 564-572.
OCTOBER 1964.
61. HISER, H.W., SENN H.V. AND DAVIES, P.J.
MESO-SCALE SYNOPTIC ANALYSIS OF RADAR AND SATELLITE
METEOROLOGICAL DATA.
MIAMI UNIV., CORAL GABLES, FLA., INST. OF MARINE
SCIENCE, CONTRACT CWB 10242, FINAL REPORT, OCT.
1963.
62. HISER, HOMER W., FRESEMAN, WILLIAM L.
RADAR METEOROLOGY.
2ND ED. REVISED BY HISER. CORAL GABLES, FLA.
UNIV OF MIAMI. 1959. 267P
63. HOWELL, WALLACE E.
THE STEREO NEPHOGRAPH--A DEVICE TO MAP CLOUDSCOPES FROM
AERIAL PHOTOGRAPHS.
HOWELL WALLACE ASSOC. INC., LEXINGTON, MASS.
CONTR AF(604)-7281. OCT. 30, 1961.
64. HUFF, F.A. AND NEILL, J.C.
FREQUENCY RELATIONS FOR STORM RAINFALL IN ILLINOIS.
ILLINOIS. STATE WATER SURVEY DIV., BULLETIN 46,
1959.
65. HWANG, S.S., TANG, M.M. AND DANG, J.C. (NANKING UNIV., DEPT
OF METEOROLOGY)
A GRAPHICAL MONTH-TO-MONTH CORRELATION METHOD OF LONG
RANGE WEATHER FORECASTING.
ACTA METEOROLOGICA SINICA, PEKING. 30(4), 306-315
NOVEMBER 1959.
66. INTERNATIONAL COMMITTEE FOR THE STUDY OF CLOUDS.
INT. ATLAS OF CLOUDS AND OF TYPES OF SKIES. 44 P
1939.

67. IVES, RONALD L.
AREAS OF OCCURENCE OF GRINDSTONE CLOUDS.
WEATHERWISE, BOSTON. 11(6), 201-203. DEC. 1958.
68. JENKINS, CARL F. AND P.F. TWITCHELL
STUDY AND EVALUATION OF 2 HOUR OBJECTIVE TERMINAL
FORECAST TECHNIQUES.
FINAL REPORT AF(604)--2450, 1959. 219P
69. JOHNSON, DAVID S.
METEOROLOGICAL SATELLITES AND WEATHER FORECASTING.
VOICE OF AMER. FORUM LECTURES. SPACE SCIENCE SER
NO. 2 1963 7P.
70. JOLY, E.F.
CODES FOR IBM WEATHER CODES USED TO RECORD FIRE WEATHER
AND GENERAL CLIMATIC DATA.
CANADA. DEPT. OF FORESTRY. MIMED 62--5 1962
71. KATS, ABRAM L'VOVICH
PREDSKAZANIE POGODY NA TRI-SEM' DNEI. (WEATHER
FORECASTING FOR THREE TO SEVEN DAYS.)
LENINGRAD. GIDROMETEORIZDAT, 1958 ED. 131P.
DIST. BY U.S. OFFICE OF TECH. SERVICE, AS ITS OTS
59-13900.
72. KEIGLER, J.E. AND KRAWITZ, L.
WEATHER RADAR OBSERVATIONS FROM AN EARTH SATELLITE.
JOURN. OF GEOPHYSICAL RESEARCH WASH. D.C. 65(9),
2793-2808 SEPT. 1960.
73. KERLEY, M. J.
ANALYSIS OF CLOUD DISTRIBUTION BETWEEN LONDON AIRPORT
AND JOHANNESBURG FROM COMET IN-FLIGHT REPORTS.
MET. MAGAZINE, LONDON. 89(1060), 297-300 NOV. 1960
74. KHRABROV, IN B.
METODIKA SOSTAVLENIIA PROGNOZOV POGODY NA 3-7 DNEI.
(METHODS OF COMPILING WEATHER FORECASTS FOR 3 TO 7
DAYS.)
MOSCOW. GIDROMETEORIZDAT, 1959 182P.
75. KHRGIAN, A. KH.
O SOVETSKOM 'ATLASE OBLAKOV' I O KLASSIFIKATSII
OBLACHNYKH FORM. (THE SOVIET CLOUD ATLAS AND A CLASS-
IFICATION OF CLOUD FORMS.)
METEOROLOGIYA I GIDROLOGIYA. NO. 3, 53-56. 1960
76. KISS, E.
BIBLIOGRAPHY ON THE USE OF SATELLITES IN METEOROLOGY.
MET. AND GEOASTROPHYSICAL ABSTRACTS VOL. 14,

1963 PP. 870-936.

77. KLEIN, W.H., CROCKETT, CURTIS W. AND ANDREWS, J.F.
OBJECTIVE PREDICTION OF DAILY PPT. AND CLOUDINESS.
JOUR. OF GEOPHYSICAL RES., WASH. D.C. 70(4),
801-813. FEBRUARY 15, 1965.
78. KOLDOVSKY, MILAU
PHOTOGRAPHIC OBSERVATION OF THE DEVELOPMENT OF A
THUNDERSTORM CLOUD.
STUDIA GEOPHYSICA ET GEODAETICA. PRAGUE 3(1),93-
97. 1959.
79. KONDROT'EV, N.N.
O METODE I NEKOTORYKH REZUL'TATAKH BAZISNOI FOTOTEODOL-
ITNOI S'EMKI OBLAKOV. (METHODS AND SOME RESULTS OF
PHOTOTHEODOLITE STEREOSCOPIC SURVEY OF CLOUDS.)
TSENTRAL' NAIA AEROLOGICHESKAIA OBSERVATORIIA,
NO. 30,84-89 1959.
80. KRISHNA, RAO. P. AND WINSTON, JAY S.
INVESTIGATION OF SOME SYNOPTIC CAPABILITIES OF
ATMOSPHERIC WINDOW MEASUREMENTS FROM SAT. TIROS II.
JOUR. OF APPLIED MET., BOSTON, 2(1),12-23, FEB.
1963.
81. KRUEGER, ARTHUR F. FRITZ, SIGMUND
CELLULAR CLOUD PATTERNS REVEALED BY TIROS I.
TEELUS. VOL. 13(1) FEB. 1961. P1-7
82. LOPEZ, MANUEL E.
REPORT OF PHOTOGRAMMETRY OF CLOUDS FROM FIXED CAMERA
SITES ON THE GROUND.
HOWELL WALLACE E. ASSOC. INC. LEXINGTON, MASS.
CONTR AF19(604)-7281. SCIENTIFIC REPORT NO 2.
DECEMBER 30, 1961.
83. LORENZ, EDWARD N.
PROSPECTS FOR STATISTICAL WEATHER FORECASTING.
MIT DEPT. OF METEOROLOGY CONTR. AF19(604)/1566
(STATISTICAL FORECASTING PROJECT) FINAL REPORT
JANUARY 1959. 185P
84. LUDLAM, F.H., R.S. SCORER
CLOUD STUDY- A PICTORIAL GUIDE.
LONDON. J. MURRAY 80 P. 1957
85. LUDLAM, F.H., R.S. SCORER
FURTHER OUTLOOK (CLOUD STUDY).
LONDON. A. WINGATE 174 P. 1954

86. LYNCH, HENRY B. (CONS. ENGS. L.A., CALIF.)
RAINFALL PATTERNS ALONG THE PACIFIC COAST, VARIATIONS I
MONTHLY RAINFALL.
1954.
87. MCDONALD, J.E. (INST ATMOS PHYSICS UNIV. OF ARIZ.)
CLOUD PHOTOGRAPHY AND CLOUD PHYSICS.
WEATHER. LONDON. 14(8),247-252 AUG. 1959.
88. MACE, L.M. AND JONES, J.B.
OPERATIONAL ASSISTS BY TIROS.
WEATHERWISE, BOSTON. VOL. 15(3), 97-101. JUNE
1962.
89. MACH, R.E. AND HEMPEL, H.P.
RESEARCH ON THE PROCESSING OF SAT. PHOTOGRAPHY BY
DIGITAL TECHNIQUES.
IBM CORP., KINGSTON, N.Y. CONTRACT AF 19(604)8432
FINAL REPORT, JUNE 30, 1962, 95P. (AFCR2-TR-62-8)
90. MADVIG, R.M.
PHASE II NIMBUS DATA HANDLING SYSTEM.
SRI, MENLO PARK, CALIF. CONTRACT NAS5 1882, FINAL
REPORT. JANUARY 1963.
91. MALKUS, JOANNE STARR, RIEHL, H.
CLOUD STRUCTURE AND DISTRIBUTION OVER THE TROPICAL
PACIFIC OCEAN.
BERKELEY, U.C. PRESS 229P. 1964.
92. MARGGRAF, WALTER A.
AUTOMATIC DATA PROCESSING OF WEATHER SATELLITE DATA.
GENERAL DYNAMICS/ASTRONAUTICS, SAN DIEGO. CONTR
AF 19 (604) 8861, REPORT GDA63 0046 JAN. 31,
1963. 49P
93. MARGGRAF, WALTER A.
WEATHER SATELLITE DATA PROCESSING.
GENERAL DYNAMICS/ASTRONAUTICS, SAN DIEGO, CONTR
19(604) 8861. FINAL REPORT JAN. 1964. 92P.
94. MERRILL, HARRISON J. (US ARMY SIGNAL RES AND DEV LAB)
CLOUD COVER OBSERVATION AND RADIATIVE MEASUREMENT FROM
SATELLITES.
N.Y. ACADEMY OF SCIENCES, TRANSACTIONS SER. 2
21(1), 153-158. DEC. 1958.
95. METEOROLOGICAL AND GEOASTROPHYSICAL ABSTRACTS
SPECIAL BIBLIO ON THE USE OF SATELLITES IN METEOROLOGY.
VOL. 15(2 AND 3). AMERICAN METEOROLOGICAL SOC.

96. MILLER, L.I.
PROCESSING SATELLITE WEATHER DATA--PART II.
1962 FALL JOINT CONF. AFIPS CONF. VOL. 22, P19-26
97. MILLER, ROBT. G.
STATISTICAL PREDICTIONS BY DISCRIMINANT ANALYSIS.
TECHNICAL MEMORANDUM NO. 4, CONTRACT NO. AF30 635
14459, HARTFORD CONN., TRAVELERS RESEARCH CENTER
INC. 138PP. 1961
98. MILLER, ROBERT R.
WEATHER BUREAU AUTOMATIC PICTURE TRANSMISSION GROUND
STATION ENGINEERING MODEL.
US WEATHER BUREAU. INSTRUMENTAL ENG. DIV.
REPORT NO 6. MAY 1964.
99. MIT
EXTENDED FORECASTING RESEARCH PROJECT.
CONTR. CWB 9232 FINAL RPT. JULY 1, 1957 JUNE 30,
1958.
100. MORRIS, WILLIAM G.
STATISTICAL ANALYSIS OF FIRE WEATHER STATION
DISTRIBUTION.
J. FORESTRY. V38 APRIL 1940 P.318-321
101. MILLER, WALTER
METHODIN ZUR ABSCHATZUNG DER AUTEILE BESTIMMTER BEWOLK
UNGSTUFEN ANS DEN DREI KLIMATERMINEN. (METHODS FOR
ESTIMATING THE FREQUENCIES OF SPECIFIED CLOUD AMOUNTS
ON THE BASIS OF THREE STANDARD OBSERVATIONS.)
WELTER UND LEBEN, VIENNA. 12(3/4), 72-76. 1960.
102. NAGLE, ROLAND E. AND BLACKMER, ROY H. JR.
USE OF SYNOPTIC SCALE WEATHER RADAR OBSERVATIONS IN THE
INTERPRETATION OF SATELLITE CLOUD OBSERVATIONS.
SRI. MENLO PARK, CALIF. CONTRACT AF 19(628) 284
FINAL REPORT, DEC. 1962. 255P. (AFCLR 63-223)
103. NORDBERG, W., BANDEEN, W.R., CONRATH, B.J. ET AL
PRELIMINARY RESULTS OF RADIATION MEAS. FROM TIROS III
JOUR. OF ATMOSPHERIC SCIENCES, VOL. 19(1), 20-29
1962.
104. NORDBERG, WILLIAM
RESEARCH WITH TIROS RADIATION MEASUREMENTS.
ASTRONAUTICS AND AEROSPACE ENGIN. N.Y. 1(3),
76-83. APRIL 1963.
105. NORDO, J.
EXPECTED SKILL OF LONG-RANGE FORECASTS WHEN DERIVED

FROM DAILY FORECASTS AND PAST WEATHER DATA.
NORWAY. METEOROLOGISHE INSTITUTT. SCIENTIFIC
REPORT NO 4. 1959.

106. O'CONNOR, JAMES F.
CLOUD SIGNIFICANCE AND FORECASTING.
AMER. METEOROLOGICAL SOCIETY, BULLETIN 34(8),
339-350. OCTOBER 1953.
107. OLIVER, V.J. AND OLIVER, M.B. (USWB WASH. D.C.)
CLOUD PATTERNS.
MONTHLY WEATHER REVIEW WASH. D.C. 91(10/12),
621-629. OCTOBER/DECEMBER 1963.
108. OSTBY, FREDERICK P. JR., CHISSHOLM, DONALD A., BALL, ROBERT
RELATIONS DERIVED BETWEEN CIRCULATION AND TOTAL CLOUD
AMOUNT BY STATISTICAL METHODS.
TRAVELERS RESEARCH CENTER, INC. HARTFORD, CONN.
CONTR AF 19(626)-16. TECH MEMO 4-7045-95. SEPT.
109. PARSONS, JAMES J. (UC BERKELEY)
FOG DRIP FROM COASTAL STRATUS, WITH SPECIAL REFERENCE
TO CALIFORNIA.
WEATHER. LONDON 15(2), 58-62 FEB. 1960.
110. PED' D.A. AND TURKETTI, Z.L.
UCHET KLIMATOLOGICHESKIKH DANNYKH PRI USTENKE PROGNOZOR
POGODY (CONSIDERATION OF CLIMATOLOGICAL DATA IN
EVALUATING WEATHER FORECASTS.)
METEOROLOGIYA I GIDROLOGIYA Leningrad, NO.5
40-43. 1959.
111. PIKE, ARTHUR C.
ESTIMATING THE EFFECT OF CLOUDINESS ON INCOMING SOLAR
RADIATION.
COLO. ST. UNIV. FT. COLLINS. DEPT OF ATMOSPHERIC
SCIENCE. TECH PAPER NO. 31. MAY 1962. 9P
112. PRABHAKARA, C. AND S.I. RASOOL
EVALUATION OF TIROS IR DATA.
PROC. 1ST INT. SYMP. ON ROCKET AND SATELLITE MET.
AMSTERDAM, HOLLAND 234-236 P. 1963.
113. PYLE, ROBERT L., AND MERCER, DONALD M.
CATALOGUE OF METEOROLOGICAL SATELLITE DATA TIROS I
TELEVISION CLOUD PHOTOGRAPHY
KEY TO METEOROLOGICAL RECORDS DOCUMENTATION NO.
5.31, U.S. DEPT OF COMMERCE, WEATHER BUREAU.
1961.
(TIROS II NO. 5.32)
(TIROS III NO. 5.33)

(TIROS IV NO. 5.34)
(TIROS V NO. 5.35)
(TIROS VI NO. 5.36)
(TIROS VII NO. 5.37)
(TIROS VIII NO. 5.38)

114. PYLE, ROBERT L.
DOCUMENTATION FOR TIROS IV TV DATA.
US WEATHER BUREAU. MSL REPORT NO. 16. MAY 1963.
115. RASOOL, S.I.
CLOUD HEIGHTS AND NIGHTTIME CLOUD COVER FROM TIROS
RADIATION DATA.
JOUR. OF ATMOSPHERIC SCIENCES. 21(2), 152-156.
1964.
116. ROSENDAL, HANS E.
AUTOMATIC PICTURE TRANSMISSION FROM WEATHER SATELLITES.
MARINERS WEATHER LOG. WASH. D.C. 7(5), 171-172.
SEPTEMBER 1963.
117. RUZECKI, MARY A.
USE OF SATELLITE CLOUD PHOTOGRAPHS IN NUMERICAL WEATHER
PREDICTION.
US WEATHER BUREAU. MET. SAT. LAB. RPT. NO. 23.
DECEMBER 1963.
118. SANDERS, FREDERICK
THE EVALUATION OF SUBJECTIVE PROBABILITY FORECASTS.
AF CAMBRIDGE RESEARCH CENTER. TN-58-465. MIT 65P
1958.
119. SCHLOSS, MILTON (GM CORP. DEF. SYSTEMS DIV. WARREN, MICH)
CLOUD COVER OF THE SOVIET UNION.
GEOGRAPHICAL REVIEW N.Y. 52(3), 389-399 JULY
1962.
120. SCHULZ, H.
ZUR ANSWERTUNG VON WOLKENMESSBILDERN MIT ELEKTRONISCHEN
RECHENAUTOMATEN. (EVALUATION OF CLOUD PHOTOGRAPHS WITH
ELECTRONIC COMPUTER.)
BEITRAGE ZUR PHYSIK DER ATMOSPHERE, FRANKFURT.
A.M. 32(1/2), 78-83. 1959.
121. SCORER, R.S.
DYNAMICS OF CLOUD FORMATIONS.
INT'L SYMP. ON ROCKET AND SATELLITE METEOROLOGY,
1ST. WASHINGTON, D.C. 1962. PROCEEDINGS.
ED. BY H. WEXLER AND T.E. CASKEY JR. AMSTERDAM,
NORTH HOLLAND. PUBL. CO. 1963.

122. SELLERS, WILLIAM D.
THE ANNUAL AND DIURNAL VARIATIONS OF CLOUD AMOUNTS AND
CLOUD TYPES AT SIX ARIZONA CITIES.
U. OF ARIZONA INST. OF ATMOSPHERIC PHYSICS,
SCIENTIFIC REPORT NO 8 1958. 104P
123. SEREBRENY, S.M. AND BLACKNER, R.H. JR.
CLOUD RESEARCH DATA.
SRI, MENLO PARK, CALIF. CONTRACT AF 19(604) 7312
FINAL REPORT, FEBRUARY 1963.
124. SEREBRENY, SIDNEY M. AND BLACKNER, RUY H. JR.
AN INVESTIGATION TO ESTABLISH THE TRUE NATURE OF CLOUD
COVER.
STANFORD RESEARCH INST. MENLO PARK, CALIF.
CONTR AF 19(604)-7312. SCIENTIFIC REPT. NO 2,
JUNE 1961. 64P
125. SEREBRENY, S.M., WIEGMAN, ELDON J. AND HADFIELD, REX G.
INVESTIGATION OF THE OPERATIONAL USE OF CLOUD PHOTO-
GRAPHS FROM WEATHER SATELLITES IN THE NORTH PACIFIC.
SRI, MENLO PARK, CALIF. CONTRACT CWB 10238,
FINAL REPORT. NOVEMBER 1962. 93P
126. SHARP, F.A.
CLOUD OVER THE OPEN SEA.
METEOROLOGICAL MAGAZINE, LONDON. 87(1033),202-204
JULY 1958.
127. SHINOHARA, T.
ERRORS ESTIMATING TEMPERATURES FROM NEIGHBORING
OBSERVATIONS.
J. OF METEOROLOGICAL RES. TOKYO. 10(2),83-87.
1958.
128. SHORR, BERNARD (ED)
STUDIES IN STATISTICAL WEATHER PREDICTION.
TRAVELER WEATHER RESEARCH CENTER. CONTRACT AF 19
(604)-1590. FINAL REPT. DECEMBER 31, 1958. 238P
129. SINGER, S.FRED
FOREST FIRE DETECTION FROM SATELLITES.
JOUR. OF FORESTRY, WASH. D.C. 61(12),860-862.
DECEMBER 1, 1962.
130. SOLOMON, IRVING
ESTIMATES OF ALTITUDE AND PROBABILITIES OF BEING ABOVE
THE CLOUDS.
U.S. AIR WEATHER SOURCE, CLIMATIC CENTER. TECH.
MEMO TO THE DIRECTOR. TM-59-2 AUG. 11, 1959.

131. SOLOMON, IRVING
ESTIMATED FREQUENCIES OF SPECIFIED CLOUD AMOUNTS WITHIN
SPECIFIED RANGES OF ALTITUDE.
U.S. AIR WEATHER SERVICE. TECH. REPORT. 167,
APRIL 1963.
132. STARK, N.
33 YR. SUMMARY OF CLIMATOLOGICAL MEASUREMENTS FROM THE
CENTRAL SIERRA NEVADA OF CALIFORNIA.
USFS PSWF AND RES. RES. NOTE. PSW 36. 1963
133. STERNBERG, SIDNEY AND WILLIAM G. STROUD
TIROS I METEOROLOGICAL SATELLITE.
ASTRONAUTICS. 5(6), 32-34 AND 84-86. JUNE 1960.
134. STRAUCH, EDWARD (PIHM, WARSAW)
FIZYCZNE PODSTAWY STOSOWANIA FILTRÓW W FOTOGRAFII CHMUR
(PHYSICAL FOUNDATIONS OF THE APPLICATION OF FILTERS IN
CLOUD PHOTOGRAPHY).
PRZEGŁÓD GEOFIZYCZNY WARSAW 5(1), 41-48. 1960.
135. STROUD, W.G.
INITIAL RESULTS OF THE TIROS I MET. SATELLITE.
J. OF GEOPHYSICAL RES. 65, 1643-1644. MAY 1960.
136. SUTTON, SIR OLIVER GRAHAM
LOOKING AHEAD IN WEATHER FORECASTING.
GEOGRAPHICAL MAGAZINE, LONDON 33(5), 287-294
SEPTEMBER 1960.
137. TOMKIN, RUSSELL J., LA RUE, J.A., SANDERS, F.
OBJECTIVE PREDICTION OF CLOUDS AND PRECIPITATION USING
VERTICALLY INTEGRATED MOISTURE AND ADIABATIC VERTICAL
MOTIONS.
J. OF APPLIED METEOROLOGY. 4(1), 3-17. FEB. 1965
138. TURNER, J.A.
THE EVALUATION OF RAINFALL PROBABILITY FORECASTS FOR
FORESTRY.
CANADA. METEOROLOGICAL BRANCH, CIR 3148, TEC 290
JANUARY 15, 1959.
139. U.S. CIVIL AERONAUTICS ADMINISTRATION
PILOT'S WEATHER HANDBOOK.
CAA TECH. MANUAL NO. 104, REV. 1955 WASH. D.C
140. U.S. METEOROLOGICAL SATELLITE LAB. SUTLAND, MD.
U.S. GODDARD SPACE FLIGHT CENTER, GREENBELT
TIROS II RADIATION DATA USERS' MANUAL.
AERONOMY AND MET. DIV., GODDARD SPACE FLIGHT CTR,
NASA AND MSL. 1961. 57PP.

141. U.S. NAVY RESEARCH FACILITY
VERTICAL MOTION AND WEATHER. (REPORT ON TASK 30--EVAL-
UATION OF NUMERICAL WEATHER PREDICTION TECHNIQUES.)
NWRF 30-0359-024. MARCH 1959.
142. U.S. WEATHER BUREAU
CLIMATOLOGICAL DATA CALIFORNIA.
US WEATHER BUREAU.
143. U.S. WEATHER BUREAU
THE CLIMATE OF HUMBOLDT--DEL NORTE CO.
EUREKA, CALIF. FARM ADVISOR. AGRIC. EXTENSION
SERVICE, U.C. 1964. 34P
144. U.S. WEATHER BUREAU
CLOUD CODE CHART.
REV. JAN. 1958
145. U.S. WEATHER BUREAU
INT'L METEOROLOGICAL SATELLITE WORKSHOP.
INT. METEOROLOGICAL SATELLITE WORKSHOP, WASH,
D.C. 1961. PROCEEDINGS.
146. U.S. WEATHER BUREAU
KEY TO METEOROLOGICAL RECORDS DOCUMENTATION.
NO. 3.151
147. U.S. WEATHER BUREAU
LOCAL CLIMATOLOGICAL DATA.
US WEATHER BUREAU.
148. U.S. WEATHER BUREAU
SYNOPTIC METEOROLOGY AS PRACTICED BY THE NATIONAL
METEOROLOGICAL CENTER, PT. 1--ANALYSIS.
THE NAWAC MANUAL, BY THE STAFF, NAT'L WEATHER
ANALYSIS CENTER WASHINGTON, D.C. OCT. 1960.
149. U.S. WEATHER BUREAU
WORLD WEATHER RECORDS, WASHINGTON D.C.
USWB WASHINGTON D.C. 1920-
150. VACHON, D.N. AND KING, J.
HIGH ALTITUDE CLOUD PHOTOGRAPHY FROM BALLISTIC MISSILES
PHILA., PENN., MISSILE AND SPACE VEHICLE DEPT.
GENERAL ELECTRIC CO. (1960 *)
151. VISHER, STEPHEN S.
CLIMATIC ATLAS OF THE UNITED STATES.
CAMBRIDGE, MASS. HARVARD U. PRESS, 1954

152. WARK DAVID O. AND WINSTON, JAY S.
APPLICATION OF SATELLITE RADIATION MEASUREMENTS TO
SYNOPTIC ANALYSIS AND TO STUDIES OF THE PLANETARY HEAT
BUDGET.
INT. SYMP. ON ROCKET AND SAT. MET. 1ST, WASH.
D.C. 1962. PROCEEDINGS. ED. BY H. WEXLER AND
T.E. CASKEY JR. AMSTERDAM, NORTH HOLLAND. PUBL.
CO. 1963. P247-253.
153. WARK, D.O., YAMAMOTO, LIENESCH, J.H.
METHODS OF ESTIMATING INFRARED FLUX AND SURFACE
TEMPERATURE FROM METEOROLOGICAL SATELLITES.
J. OF ATMOSPHERIC SCIENCES. 19(5). SEPT. 1962.
154. WEINSTEIN, MELVIN AND SUOMI, VERNER E.
ANALYSIS OF SATELLITE I-R RADIATION MEASUREMENTS ON
A SYNOPTIC SCALE.
MONTHLY WEATHER REVIEW. VOL. 89(11), 419-428.
NOVEMBER 1961.
155. WEXLER, HARRY AND FRITZ, SIGMUND (USWB)
TIROS REVEALS CLOUD FORMATIONS.
SCIENCE, WASH. D.C. 131(3415), 1708-1710.
JUNE 10, 1960.
156. WHITE, ROBERT M., MYERS, R.F., WARD, F.W. AND GUY, G.A.
GENERAL DESIGN FOR WEATHER OBSERVING AND FORECASTING
SYSTEM.
USAF CAMBRIDGE RES. CENTER AFRCR TN 57 609.
ASTIA DOCUMENT 133846. DEC. 15, 1957. 92P
157. WIDGER, WILLIAM K. JR. AND WOOD, CHAS.P.
EXPLANATION OF THE LIMITATIONS TO THE COVERAGE PROVIDED
BY TIROS.
WEATHERWISE, BOSTON. 14(6), 230-237. DEC. 1961.
158. WIDGER, WILLIAM K.
METEOROLOGICAL SATELLITES AND WEATHER RECONNAISSANCE
AIRCRAFT--COMPLEMENTARY OBSERVING SYSTEMS.
AMERICAN METEOROLOGICAL SOCIETY, BULLETIN
44(9), 549-563. SEPT. 1963.
159. WINSTON, JAY S. AND TOURVILLE, LLOYD (USWB)
CLOUD STRUCTURE OF AN OCCLUDED CYCLONE OVER THE GULF OF
ALASKA AS VIEWED BY TIROS I.
AMERICAN METEOROLOGICAL SOCIETY, BULLETIN.
42(3), 151-165. MARCH 1961.
160. WINSTON, J.S.
THE OPERATIONAL USE OF METEOROLOGICAL SATELLITE DATA.
ANNALS OF THE N.Y. ACAD. OF SCIENCES V93/19

OCT. 3, 1962. P775-812

- 161. WORLD METEOROLOGICAL ORG.
INT. CLOUD ALBUM FOR OBERVERS IN AIRCRAFT.
GENEVA. 1956
- 162. WORLD METEOROLOGICAL ORG.
INT. CLOUD ATLAS.
GENEVA. 1956.
- 163. YAMAGUCHI, RANOU (AVIATION WEATHER SERVICE, TOKYO)
CLOUD DISTRIBUTION REVEALED BY TIROS III.
METEOROLOGICAL SOCIETY OF JAPAN, TOKYO. JOUR.
SER. 2 42(5), 285-298. OCT. 1964.